

( )

**(*Arachis hypogaea* L.)**

\*

( / / : // : )

( )

( / / / )

( )

/

( )

)

(

/ /

:

.et al., 1997)

(Bell et al., 1987; Bell et al.,

.1993)

(Wilcult & Richburg, 1994; Rehman et al.,

.2001)

.(Bridges et al., 1992)

(Tahbaz & Sane Shariat

Panahi, 1992; Kochaki et al., 1994; Andriw &  
.Wilcut, 2002)

.(Bell et al., 1993; Barry et al., 1996)

(Bull et al., 1987; Tahbaz & Sane Shariat

.Panahi, 1992)

.(Choudhury et al., 1997)

(Kochaki et al., 1994; Rashed Mohassel et al.,

.1996; Andriw & Wilcut, 2002)

( )

.(Ghosh & Biswas, 1995)

.(Patel et al., 1985)

(Kasai et al., 1997)

(Jaffar & Gardner, 1988; Fiebig et al., 1991)

(Hauser & Buchanan, 1981; Jaffar &

.Gardner, 1988; Kasai et al., 1997)

.(Bridges et al., 1992)

(Bridges et al.,

.1992)

(Raghvani et

(Gardner & Auma, 1989; Barry et al., 1996; Kasai



(*Amaranthus retroflexus* L.)  
(*Portulaca* (*Chenopodium album* L.)  
(*Solanum nigrum* L.) *oleracea* L.)

(*Cynodon dactylon* (*Artemisia biennis* Willd.)  
(*Digitaria sanguinalis* L.) L.)  
(*Datura* (*Eleusine indica* L.)  
*stramonium* L.)

(*Daucus carota* L.)  
(*Paspalum* (*Veronica persica* poir)  
(*Cyperus esculantus* *dilatatum* poir)  
L.)

(Bell et al., 1993)

( )

)

(

)

(

(Barry et al., 1996)

( )

(Bell et al., 1993)

( )

SAS

EXCEL

)

(

...

:

(Jaffar & Gardner, 1988; Fiebig et al., 1991)

(Andriw & Wilcut, 2002)

(Fiebig et al., 1991)

)

(

(David et al., 2001)

(Hausar et al., / (Choudhury et al., 1997)

(Fiebig et al., 1991) / 1979)

(Ghosh & Biswas, 1995)

(Barry et al., 1996)

(Andriw & Wilcut, 2002)

/

( )

)

(Gardner & Auma, 1989; David

(

.et al., 2001)

(Choudhury et al., 1997; David et al.,

.(Raghvani et al., 1984)

.2001)

/

( )

/

/	ns	/	**	/	ns	/	ns
/	ns	/	ns	/	*	/	ns
/	*	/	*	/	*	/	ns
/	ns	/	ns	/	ns	/	ns
/		/		/		/	
/		/		/		/	(%)

\* \*\* ns

.( )

.(Hausar et al., 1979)

( )

...

:

/ b / b / b / b / b / b	/ b / b / b / b / b / b	/ b / b / b / b / b / b
/ a / a / a / a / a / a	/ a / a / a / a / a / a	/ a / a / a / a / a / a
a / a / a / a / a / a	/ a / a / a / a / a / a	/ a / a / a / a / a / a
/ b / b / b / b / b / b	/ b / b / b / b / ab / ab	/ b / b / b / ab / ab / ab
/ bc / c / b / bc / c / c	/ c / c / c / bc / bc / bc	/ c / b / b / bc / bc / bc
/ c / d / c / c / d / d	/ c / d / d / c / c / c	/ d / c / c / c / c / c

.( )

/ ns	/ ns	/ **
/ ns	/ ns	/ *
/ **	/ **	/ **
/ ns	/ ns	/ ns
/	/	/
/	/	/ (%)

\* \*\* ns

(Barry et al., 1996)

)

( )

(

)

(

(Jaffar & Gardner, 1988;

.Munda & Patel, 1989)

/ / / (%)

\* \*\* ns

/ \* / \* / \*  
/ \* / \* / \*\*  
/ \*\* / \*\* / \*\*  
/ ns / ns / ns  
/ / /  
/ / / (%)



\* \*\* ns

/ a / a / a	/ a / a / a
/ b / b / b	/ b / b / b
/ a / a / a	/ a / a / a
/ b / b / b	/ b / b / b /
/ c / c / c	/ c / c / b /
/ d / d / d	/ d / d / c /

.( )

/ \* / \*\* / \*  
/ \* / \*\* / \*  
/ \*\* / \*\* / \*\*  
/ ns / ns / ns  
/ / /





( ) .

## REFERENCES

- Andriw, J. & Wilcut, J. W. (2002). Weed management with diclosulam in Strip- Tillage peanut (*Arachis hypogaea* L.). *Weed Technology*, 16, 29- 36.
- Barry, J. B., Funderburk, J. E., Teare, I. D. & Gorbet, D. W. (1996). Interaction of early –season herbicide hnjury, tobacco trips injury and cultivar on peanut. *Agron J*, 88, 14- 18.
- Bell, M. J., Muchow, R. C. & Wilson, G. L. (1987). The effect of plant population on peanut (*Arachis hypogaea* L.) in a monsoonal tropical environment. *Field Crop Research*, 17, 91- 107.
- Bell, M. J., Wright, G. C. & Harch, G. R. (1993). Enviromental and agronomic effects on the growth of four peanut cultivars in a sub- tropical enviroment. I. Dry matter acumulation and radiation use efficiency. *Experimental Agriculture*, 29 (4), 473- 490.
- Bridges, D. C., Brecke, B. J. & Barbour, J. C. (1992). Wild poinsettia (*Euphorbia heterophylla*) interference with peanut. *Weed Sci*, 40 (1), 37-42.
- Choudhury, B., Zahid, M., Ghosh, P. & Samui, R. C. (1997). Effect of rate and row spacing on confectionary groundnut in West Bengal, India. *International Arachis Newsletter*, 17, 71- 73.
- David, L., Jordan, B., Beam, P., Dewayne, J. & Spears, F. (2001). Peanut response to prohexadione calsium in three seeding rate, row pattern planting system. *Agron J*, 93, 232- 236.
- Fiebig, W. W., Shilling, D. G. & Knauft, D. A. (1991). Peanut genotype response to interference from common cocklebur. *Crop Sci*, 31(5), 1289- 1292.
- Gardner, F. P. & Auma, E. Q. (1989). Canopy structure, light interception and yield market quality of peanut genotypes as influenced by planting pattern and planting date. *Field Crop Research*, 20 (1) , 13- 29.
- Ghosh, A. K. & Biswas, A. K. (1995). Regulation of corelative senesence in *Arachis hypogaea* L. by source – sink aleration through physical and hormonal means. *Agron J*, 175, 195- 202.
- Gill, M. A., Islam, J. S., Brar, H. S. & Mehra, S. P. (1986). Studies on the critical period of weed competition in groundnut. *Weed Abstracts*, 37 (5), 195.

12. Hausar, E. W., Buchanum, G. A. & Slaughter, J. W. (1979). Effects of row spacing, weed- free maintenance periods and herbicide systems on the yield of florunner peanuts. *Proceedings American Peanut Research and Education Society*, 11 (1), 42.
13. Hauser, E. W. & Buchanan, G. A. (1981). Influence if row spacing , seeding rates, and herbicide system on competativeness and yield of peanut. *Peanut Sci*, 8, 74- 81.
14. Jaffar, Z. & Gardner, F. D. (1988). Canopy development, yield and market quality in peanut as affected by genotype and planting pattern. *Crop Sci*, 28, 299- 305.
15. Kasai, F. S., Paulo, E. M. & Cavichioli, J. C. (1997). Effects of weed competition periods on peanut crops: Dry season peanut of 1988. *Bragantia*, 56 (2), 323- 331.
16. Kochaki, A., Hoseini, M., Nasiri Mahallati, M. & Khiyabani, H. (1994). *Weed Ecology*. Tehran University Press. (In Farsi).
17. Munda, D. C. & Patel, C. S. (1989). Data of sowing, spacing and nutrient requirement of groundnut (*Arachis hypogaea* L.) under mid- altitude of meghalaya. *Indian Journal of Agricultural Sci*, 59 (11), 706- 709.
18. Patel, R. G., Patel, R. B. & Patel, T. D. (1985). Crop weed competition in summer groundnut. *Weed Abstracts*, 35 (6), 234.
19. Raghvani, B. R., Goyal, S. N. & Patel, J. C. (1984). Crop weed competition in rainfed groundnut(*Arachis hypogaea* L.) . *Indian Journal of Weed Sci*, 16 (4), 234- 237.
20. RAO, V. S. (1992). *Principles of Weed Science*. Oxford & IBH publishing co. PVT. LTD.
21. Rashed Mohassel, M. H., Rahimian, H. & Banayan, M. (1996). *Applied Weed Science*. Jahad University of Mashhad Press. 5760p. (In Farsi).
22. Rehman, A. U., Wells, R. & Isleib, T. G. (2001). Reproductive allocation on branches of virinia tape peanut cultivars bred for yield in north carolina. *Crop Sci*, 41, 72- 77.
23. Tahbaz, F. & Sane Shariat Panahi, M. (1992). *Weed Biology*. Tehran University Press. 2700p. (In Farsi).
24. Walker, R. H., Wells, L. W. & Guire, M. C. (1989). Bristly starbur (*Acanthospermum hispidum*) interference in peanuts (*Arachis hypogaea* L.). *Weed Sci*, 37(2), 196- 200.
25. Widaryanto, E. (1994). Influence of preemergence doses of the herbicide oxyfelfuorfen (Goal 2E) and population density of peanut (*Arachis hypogaea* L.) on weed control and its effect on growth and yield of peanut plants on dry land. *Procceding American Peanut Research and Education Society*. 12, 36.
26. Wilcult, J. W. & Richburg, J. S. (1994). Imazethapyre and paraquat systems for weed management in peanut (*Arachis hypogaea* L.). *Weed Sci*, 42, 601- 607.